

UNCLASSIFIED

---

AD **260 729**

*Reproduced  
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA



---

UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGED BY ASTIA  
AS AD NO.

260729

# DUALITY IN HOMOGENEOUS PROGRAMMING

by

E. Eisenberg

RESEARCH REPORT 8

26 JUNE 1961


I.E.R. 172-9

OPERATIONS RESEARCH CENTER

INSTITUTE OF ENGINEERING RESEARCH

XEROX

UNIVERSITY OF CALIFORNIA - BERKELEY



# DUALITY IN HOMOGENEOUS PROGRAMMING

by

E. Eisenberg  
Operations Research Center  
University of California, Berkeley

26 June 1961

Research Report 8

Work on this paper was supported, in part, by the Logistics Branch of the Office of Naval Research under Contract Nonr-562(15) at Brown University and the Bureau of Supplies and Accounts of the Office of Naval Research under Contract Nonr-222(83) at the University of California, Berkeley. Reproduction in whole or in part is permitted for any purpose of the United States Government.

## DUALITY IN HOMOGENEOUS PROGRAMMING

The problem of maximizing a concave function subject to linear constraints does not have a dual, as is the case in linear programming, in which primal optimizing variables do not appear. As a special case of our principal result it will follow that such a dual does indeed exist whenever the objective function is also homogeneous.

In the linear case we are given an  $m \times n$  matrix  $A$  and vectors<sup>\*</sup>  $a \in R^n$ ,  $b \in R^m$ . The feasibility sets  $X$  and  $Y$  are defined by:  $X = R_+^m \cap \{x | xA \leq a\}$ ,  $Y = R_+^n \cap \{y | Ay \geq b\}$ . Since  $xA \leq a$  if and only if  $xAy \leq ay$  for all  $y \in R_+^n$  (and similarly for  $Ay \geq b$ ), we may write:

$$(1) \quad \begin{aligned} X &= R_+^m \cap \left\{ x \mid xAy \leq \psi(y) \quad \text{all } y \in R_+^n \right\} \\ Y &= R_+^n \cap \left\{ y \mid xAy \geq \phi(x) \quad \text{all } x \in R_+^m \right\} \end{aligned}$$

where  $\psi(y) = ay$  and  $\phi(x) = bx$ .

A fundamental theorem of linear programming (see, e.g., [3] and [5]) states that if  $X$  and  $Y$  are both non-empty then

$$(2) \quad \max_{x \in X} \phi(x), \min_{y \in Y} \psi(y) \text{ exist and are equal.}$$

We propose to demonstrate that (2) holds for larger class of triples  $(A, \phi, \psi)$ .

---

\*  $R^m$  denotes the set of all real  $m$ -tuples. If  $u, v \in R^m$  then  $u \leq v$  means that the inequality holds for each component. In particular,  $R_+^m = R^m \cap \{x \mid x \geq 0\}$ . If  $M$  is a  $p \times q$  matrix and  $N$  is a  $q \times t$  matrix then  $MN$  represents the usual matrix product. To simplify notation, the same symbol is used for both a column vector and its transpose; the meaning will, in any case, be clear from the context.

Assumption  $A_1$ . Let  $\phi : R_+^m \rightarrow R$ ,  $\psi : R_+^n \rightarrow R$  be positively homogeneous\*, continuous, concave and convex respectively.

Let us first show that  $A_1$  does not guarantee that (2) holds when  $X$  and  $Y$  are non-empty. If  $m = 2$ ,  $n = 1$  and  $A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ ,  $\phi(x) = \phi(\xi, \eta) = \frac{\xi \eta}{\xi + \eta}$ ,  $[\phi(0) = 0]$ ,  $\psi(y) = y$  then  $A_1$  is satisfied and  $X = R_+^2 \cap \{(\xi, \eta) \mid \eta \leq 1\}$ ,  $Y = R_+ \cap \{y \mid y \geq 1\}$  are non-empty. Thus  $\min_{y \in Y} \psi(y) = 1$ , but if  $\eta \leq 1$  then  $\phi(\xi, \eta) < 1$ , although  $\sup_{x \in X} \phi(x) = 1$ , hence  $\max_{x \in X} \phi(x)$  does not exist.

The situation just illustrated cannot occur if the following holds:

Assumption  $A_2$ .

- i) If  $x \in R_+^m$ ,  $xA \leq 0$ ,  $\phi(x) \geq 0$  then  $x = 0$
- ii) If  $y \in R_+^n$ ,  $Ay \geq 0$ ,  $\psi(y) \leq 0$  then  $y = 0$

One sees immediately that (i) is violated in the preceding example, for let  $x = (1, 0)$  then  $xA = 0$  and  $\phi(x) = 0$ .

Before proving that if  $A_1$  and  $A_2$  hold then so does (2), we require the following lemma which specializes to homogeneous functions the well-known fact that a concave function is the infimum of its supports. The proof is presented here for the sake of completeness.

Lemma 1. Let  $\phi$  be as in assumption  $A_1$ , consider  $T = R^m \cap \{t \mid tx \geq \phi(x) \text{ all } x \in R_+^m\}$ , then  $T$  is non-empty, and  $\phi(x) = \inf_{t \in T} tx$ , for all  $x \in R_+^m$ .

Proof. Let  $C = \{(x, \lambda) \mid x \in R_+^m, \lambda \leq \phi(x)\}$  then  $C$  is a closed convex cone. Now if  $x_0 \in R_+^m$ ,  $\mu > 0$ , then  $(x_0, \mu + \phi(x_0)) \notin C$ , whence (see [2], Theorem 1)

\* A function  $f : C \rightarrow R^q$ , where  $C \subset R^p$  is a cone, is positively homogeneous providing  $f(\lambda x) = \lambda f(x)$  for all  $x \in C$  and  $\lambda \in R_+$ .

there exist  $t \in R^m$  and  $\alpha \in R$  such that  $tx_0 - \alpha[\mu + \phi(x_0)] < 0 \leq tx - \alpha\lambda$  all  $(x, \lambda) \in C$ .

It then follows that  $\alpha > 0$ , so that (dividing by  $\alpha$ ) we may assume  $\alpha = 1$ , but then  $t \in T$ . Reiterating, if  $x_0 \in R_+^m$ ,  $\mu > 0$  then  $\exists t \in T$  such that:

$$tx_0 - \mu \leq \phi(x_0) \leq tx_0$$

giving the desired result. We are now able to prove:

Theorem 1. If assumptions  $A_1$  and  $A_2$  hold then (2) holds.

Proof. Let

$$(3) \quad \begin{aligned} S &= R^n \cap \left\{ s \mid sy \leq \psi(y), \quad \text{all } y \in R_+^n \right\} \\ T &= R^m \cap \left\{ t \mid tx \geq \phi(x), \quad \text{all } x \in R_+^m \right\} \end{aligned}$$

Then  $S$  and  $T$  are convex sets; now consider the system of inequalities:

$$(4) \quad \begin{aligned} (x, y, s, t) &\in R_+^m \times R_+^n \times S \times T \\ s - xA &> 0 \\ -t + Ay &> 0 \\ \phi(x) - \psi(y) &> 0 \end{aligned}$$

If (4) has a solution  $x, y, s, t$  then

$$\psi(y) < \phi(x) \leq tx \leq xAy \leq sy \leq \psi(y)$$

which is a contradiction. Thus (see [2], Theorem 1) there exist  $x_0 \in R_+^m$ ,  $y_0 \in R_+^n$ ,  $\lambda \in R_+$ , not all zero and such that  $(s - xA)y_0 + x_0(Ay - t) + \lambda[\phi(x) - \psi(y)] \leq 0$  for all  $x \in R_+^m$ ,  $y \in R_+^n$ ,  $s \in S$ ,  $t \in T$ . From the homogeneity and continuity of  $\phi$  and  $\psi$  it then follows that:

$$x Ay_0 \geq \lambda \phi(x) \quad , \quad \text{all } x \in R_+^m$$

$$x_0 Ay \leq \lambda \psi(y) \quad , \quad \text{all } y \in R_+^n$$

$$s y_0 \leq t x_0 \quad , \quad \text{all } s \in S, t \in T$$

The last condition together with Lemma 1 imply:

$$\psi(y_0) \leq \phi(x_0)$$

Now if  $\lambda = 0$  then either  $x_0 \neq 0$  or  $y_0 \neq 0$  and  $Ay_0 \geq 0$ ,  $x_0 A \leq 0$ . Suppose  $x_0 \neq 0$ , if  $y_0 \neq 0$  the argument is analogous, then by  $A_2(i)$  we have  $\phi(x_0) < 0$ , whence  $\psi(y_0) < 0$  and  $y_0 \neq 0$ , contradicting  $A_2(ii)$ . Thus  $\lambda > 0$  and, dividing all inequalities by  $\lambda$ , we may assume  $\lambda = 1$ . This tells us that  $x_0 \in X$ ,  $y_0 \in Y$  and  $\phi(x_0) \leq x_0 Ay_0 \leq \psi(y_0) \leq \phi(x_0)$ . So that if  $x \in X$ ,  $y \in Y$  then

$$\phi(x) \leq x Ay_0 \leq \psi(y_0) = \phi(x_0)$$

$$\psi(y) \geq x_0 Ay \geq \phi(x_0) = \psi(y_0)$$

proving the theorem.

In case  $\phi$  and  $\psi$  are linear-homogeneous then it is true that  $\max_{x \in X} \phi(x)$  exists if and only if  $\min_{y \in Y} \psi(y)$  exists, in which case they are equal. As above, this statement is not always true under assumption  $A_1$ ; however, we show:

Theorem 2.

- I) If  $A_1$  and  $A_2(ii)$  hold, and  $\max_{x \in X} \phi(x)$  exists then (2) holds.
- II) If  $A_1$  and  $A_2(i)$  hold, and  $\min_{y \in Y} \psi(y)$  exists then (2) holds.

We prove (I), the proof of (II) is similar. Suppose that  $x_0 \in X$  and  $\phi(x_0) = \max_{x \in X} \phi(x)$ , then the system:



$$\begin{aligned}
(6) \quad & (x, s) \in R_+^m \times S \\
& s - xA > 0 \\
& \phi(x) - \phi(x_0) > 0
\end{aligned}$$

has no solution. Thus (see [2], Theorem 1) there exist  $y_0 \in R_+^n$ ,  $\lambda \in R_+$ , not both zero and such that

$sy_0 - xAy_0 + \lambda[\phi(x) - \phi(x_0)] \leq 0$  for all  $x \in R_+^m$ ,  $s \in S$ . From the homogeneity of  $\phi$  and Lemma 1 it then follows that

$$\begin{aligned}
(7) \quad & xAy_0 \geq \lambda\phi(x), \quad \text{for all } x \in R_+^m \\
& \psi(y_0) \leq \lambda\phi(x_0)
\end{aligned}$$

Now, if  $\lambda = 0$  then  $y_0 \neq 0$  and  $Ay_0 \geq 0$ ,  $\psi(y_0) \leq 0$ , contradicting  $A_2(ii)$ . It may then be assumed that  $\lambda > 0$  and, in fact, that  $\lambda = 1$  (replacing  $y_0$  by  $\lambda y_0$ ). Thus, from (7),  $y_0 \in Y$ , and for any  $y \in Y$  we have:

$$\psi(y_0) \leq \phi(x_0) \leq x_0 Ay \leq \psi(y),$$

$$\text{i.e.,} \quad \psi(y_0) = \min_{y \in Y} \psi(y) = \phi(x_0). \quad \text{q.e.d.}$$

It should be remarked that if  $A_1$  holds then (i) and (ii) of assumption  $A_2$  are equivalent to (i)' and (ii)' respectively of:

Assumption  $A_2'$ .

- i)'  $\exists y_0 \in R_+^n \ni xAy_0 > \phi(x) \quad \text{all } x \in R_+^m, x \neq 0$
- ii)'  $\exists x_0 \in R_+^m \ni x_0 Ay < \psi(y) \quad \text{all } y \in R_+^n, y \neq 0$

These in turn are equivalent to the familiar conditions that  $X$ ,  $Y$  have non-empty interiors. To see, for instance, that (i) and (i)' are equivalent it

suffices to show that (i) implies (i)' since the implication in the other direction is trivial. Assuming (i)' false, the system

$$(8) \quad (y, t) \in R_+^n \times T$$

$$Ay - t > 0$$

has no solution, whence (see [2], Theorem 1) there is an  $x \in R_+^m$ ,  $x \neq 0$ , and such that  $xAy \leq tx$  for all  $y \in R_+^n$  and  $t \in T$ . Thus  $xA \leq 0$  and (using Lemma 1)  $\phi(x) \geq 0$ , contradicting (i). To return to our remark about maximizing a concave homogeneous and continuous function  $\phi: R_+^m \rightarrow R$ , subject to the inequalities  $x \geq 0$  and  $xA \leq a$ , the dual is then: minimize  $ay$  subject to  $y \in Y$ . Conditions (i) and (ii)' become:

$$x \in R_+^m, x \neq 0, xA \leq 0, \phi(x) \geq 0 \quad \text{has no solution; and}$$

$$x \in R_+^m, xA < a \quad \text{has a solution; respectively.}$$

Also, since  $y \in Y$  providing  $y \geq 0$  and  $Ay \geq t$  for some support  $t$  of  $\phi$ , we may characterize  $Y$  by means of the gradient of  $\phi$ .

Results similar to Theorems 1 and 2 can be shown to hold under other and somewhat less restrictive assumptions; the duality theorems of linear programming then turn out to be special cases of these theorems (3, 4 and 5).

Henceforth we assume that  $A_1$  holds and consider the sets:

$$K_1 = R^{m+n+1} \cap \left\{ (\bar{x}, \bar{y}, \lambda) \mid \exists s \in S, t \in T, x \in R_+^m, y \in R_+^n \text{ and} \right.$$

$$\left. x \geq t - Ay, \bar{y} \leq s - xA, \lambda \leq \phi(x) - \psi(y) \right\},$$

$$K_2 = R^{n+1} \cap \left\{ (y, \lambda) \mid \exists s \in S, x \in R_+^m, \text{ and } \bar{y} \leq s - xA, \lambda \leq \phi(x) \right\},$$

$$K_3 = R^{m+1} \cap \left\{ (x, \lambda) \mid \exists t \in T, y \in R_+^n, \text{ and } \bar{x} \geq t - Ay, \lambda \geq \psi(y) \right\}.$$

The sets  $K_1$ ,  $K_2$ , and  $K_3$  are readily seen to be convex (because  $\psi$  and  $-\phi$  are convex); furthermore, if  $\phi$  and  $\psi$  are linear then  $K_1$ ,  $K_2$  and  $K_3$  are also closed sets. Of course any (or all) of  $K_1$ ,  $K_2$ ,  $K_3$  may be closed without either  $\phi$  or  $\psi$  being linear. Thus it is important to know the following:

Theorem 3.

If  $K_1$  is closed and  $X, Y$  are both non-empty, then (2) holds.

Theorem 4.

If  $K_3$  is closed and  $\max_{x \in X} \phi(x)$  exists, then (2) holds.

Theorem 5.

If  $K_2$  is closed and  $\min_{y \in Y} \psi(y)$  exists, then (2) holds.

Proof of Theorem 3: If the point  $(\bar{x}, \bar{y}, \lambda) = 0$  is in  $K_1$  then (2) obviously holds, suppose  $0 \notin K_1$ . Since  $K_1$  is convex and closed, there exist (see [3])  $x_0 \in R^m$ ,  $y_0 \in R^n$ ,  $\lambda_0 \in R$ ,  $\alpha \in R$ , such that:

$$0 < \alpha \leq x_0 \bar{x} - y_0 \bar{y} - \lambda_0 \lambda, \quad \text{all } (\bar{x}, \bar{y}, \lambda) \in K_1.$$

Since  $S$  and  $T$  are non-empty (see Lemma 1), and since  $(\bar{\bar{x}}, \bar{\bar{y}}, \bar{\lambda}) \in K_1$  whenever there exist  $(\bar{x}, \bar{y}, \lambda) \in K_1$  such that  $\bar{\bar{x}} \geq \bar{x}$ ,  $\bar{\bar{y}} \leq \bar{y}$  and  $\bar{\lambda} \leq \lambda$ , it follows that  $x_0 \geq 0$ ,  $y_0 \geq 0$ , and  $\lambda_0 \geq 0$ . Also,

$$0 < \alpha \leq x_0 (t - Ay) - (s - xA)y_0 - \lambda_0 [\phi(x) - \psi(y)],$$

$$\text{all } (s, t, x, y) \in S \times T \times R_+^m \times R_+^n$$

From the homogeneity of  $\phi$  and  $\psi$  and Lemma 1 it then follows that:

$$0 < \alpha \leq \phi(x_0) - \psi(y_0)$$

$$x_0 Ay \leq \lambda_0 \psi(y) \quad \text{all } y \in R_+^n$$

$$xAy_0 \geq \lambda_0 \phi(x) \quad \text{all } x \in R_+^m$$

Thus,  $\lambda_0 \psi(y_0) \geq x_0 A y_0 \geq \lambda_0 \phi(x_0) \geq \lambda_0 \alpha + \lambda_0 \psi(y_0)$ , and  $\lambda_0 = 0$ . But then  $x_0 A \leq 0$  and  $A y_0 \geq 0$ . Now  $X, Y$  we assumed non-empty, let  $x \in X, y \in Y$ . For any  $\lambda \in R_+$  we then have:

$$\begin{aligned} \phi(x) + \lambda \phi(x_0) &\leq \phi(x + \lambda x_0) \leq (x + \lambda x_0) A y \leq \\ &\leq (x + \lambda x_0) A (y + \lambda y_0) \leq x A (y + \lambda y_0) \\ &\leq \psi(y + \lambda y_0) \leq \psi(y) + \lambda \psi(y_0). \end{aligned}$$

Thus,  $\psi(y) - \phi(x) \geq \lambda [\phi(x_0) - \psi(y_0)] \geq \lambda \alpha$  for all  $\lambda \in R_+$  which contradicts  $\alpha > 0$ . Thus  $0 \in K_1$  and (2) holds. q. e. d.

Proof of Theorems 4 and 5: We prove Theorem 4, the proof of Theorem 5 is analogous. By hypothesis  $X$  is non-empty and  $\phi$  is bounded above on  $X$ , let:

$$M = \sup_{x \in X} \phi(x)$$

If  $(x, \lambda) = (0, M)$  is in  $K_3$  then, trivially, (2) holds. We show that the contrary assumption leads to a contradiction. If  $(0, M) \notin K_3$  then, as in the proof of Theorem 4, it follows from the various properties of  $K_3$  that there exist  $x_0 \in R_+^m$ ,  $\lambda_0 \in R_+$  and  $\alpha \in R$  such that:

$$\lambda_0 M < \alpha \leq x_0(t - Ay) + \lambda_0 \psi(y), \quad \text{all } (t, y) \in T \times R_+^n$$

Hence, as before,

$$x_0 A y \leq \lambda_0 \psi(y), \quad \text{all } y \in R_+^n$$

$$\text{and } \phi(x_0) > \lambda_0 M.$$

If  $\lambda_0$  is positive then  $x = \lambda^{-1} x_0 \in X$  and  $\phi(x) > M$ , contradicting the definition of  $M$ . Thus  $\lambda_0 = 0$ , and  $x_0 A \leq 0$ ,  $\phi(x_0) > 0$ ; the last contradicts the fact that  $X$  is non-empty and that  $\phi$  is bounded above on  $X$ . q.e.d.

As a final result we demonstrate that if  $\phi$  and  $\psi$  are both linear (homogeneous) then  $K_1$  is closed. That  $K_2$  and  $K_3$  are closed, under the same linearity hypothesis, follows in a similar manner.

Suppose  $\phi(x) = bx$ ,  $\psi(y) = ay$  ( $b \in R^m$ ,  $a \in R^n$ ), first note that in this case:

$$S = \left\{ s \mid s \in R^n \text{ and } s \leq a \right\}$$

$$T = \left\{ t \mid t \in R^m \text{ and } t \geq b \right\}.$$

Next, suppose we have a sequence  $(\bar{x}_k, \bar{y}_k, \lambda_k) \in K_1$  ( $k = 1, 2, \dots$ ) which converges to  $(\bar{x}, \bar{y}, \lambda) \in R^{m+n+1}$ . Thus there exist  $(s_k, t_k, x_k, y_k) \in S \times T \times R_+^m \times R_+^n$  such that:

$$\begin{aligned} \bar{x}_k &\geq t_k - Ay_k \geq b - Ay_k \\ (9) \quad \bar{y}_k &\leq s_k - x_k A \leq a - x_k A \quad k = 1, 2, \dots \\ \lambda_k &\leq bx_k - ay_k \end{aligned}$$

and

$$(10) \quad \bar{x}_k \rightarrow \bar{x}, \quad \bar{y}_k \rightarrow \bar{y}, \quad \lambda_k \rightarrow \lambda.$$

Now, suppose  $x \in R_+^m$ ,  $y \in R_+^n$ ,  $a \in R_+$  are such that  $Ay - ab \geq 0$ ,  $xA - aa \leq 0$ .

From (9) it then follows that for each  $k$  we have:

$$\begin{aligned} x\bar{x}_k &\geq bx - xAy_k \geq bx - aay_k \geq bx - abx_k + a\lambda_k \geq \\ &\geq bx + a\lambda_k - x_k Ay \geq bx + a\lambda_k + \bar{y}_k y - ay. \end{aligned}$$

i. e.,  $x(\bar{x}_k - b) + y(a - \bar{y}_k) - a\lambda_k \geq 0$ ,  $k = 1, 2, \dots$  and, by (10),  
 $x(\bar{x} - b) + y(a - \bar{y}) - a\lambda \geq 0$ .

In summary, then, the system:

$$x \in R_+^m, \quad y \in R_+^n, \quad a \in R_+$$

$$Ay - ab \geq 0, \quad xA - aa \leq 0$$

$$x(\bar{x} - b) + y(a - \bar{y}) - a\lambda < 0$$

has no solution. It follows then from the ordinary feasibility theorem for linear inequalities (see e. q. [5]) that there is an  $x \in R_+^m$  and  $y \in R_+^n$  such that:

$$(x, y) \begin{bmatrix} A & 0 & -b \\ 0 & -A^T & a \end{bmatrix} \leq (a - \bar{y}, \bar{x} - b, -\lambda),$$

i. e.,  $xA \leq a - \bar{y}$ ,  $-Ay \leq \bar{x} - b$  and  $ay - bx \leq -\lambda$ . But, as noted before,  $a \in S$  and  $b \in T$ , thus  $(\bar{x}, \bar{y}, \lambda) \in K$ , and  $K_1$  is closed.

## REFERENCES

1. Eisenberg, E., "Supports of a Convex Function," Berkeley: University of California, Operations Research Center. Research Report 5, 22 May 1961.
2. Ky Fan, I. Glicksberg, and A. J. Hoffman, "Systems of Inequalities Involving Convex Functions," Proceedings Am. Math. Soc., vol. 8, no. 3, June 1957.
3. Fenchel, W., "Convex Cones, Sets and Functions," from notes by D. W. Blacket of Lectures at Princeton University, Spring 1953.
4. Gale, D., H. W. Kuhn, and A. W. Tucker, "Linear Programming and the Theory of Games," Activity Analysis of Production and Allocation. Cowles Commission Monograph 13, New York 1951, pp. 317-329.
5. Gale, D., Theory of Linear Economic Models, McGraw-Hill, 1960.

# BASIC DISTRIBUTION LIST FOR UNCLASSIFIED TECHNICAL REPORTS

Head, Logistics and Mathematical Statistics Branch Office of Naval Research Washington 25, D. C.	Professor G. L. Bach Carnegie Institute of Tech. Planning and Control of Industrial Operations Schenley Park Pittsburgh 13, Penn.	Professor C. E. Lemke Dept. of Mathematics Rensselaer Polytechnic Institute Troy, New York
C.O., ONR Branch Office Navy No. 100 F.P.O. New York City, New York	Professor A. Charnes The Technological Institute Northwestern University Evanston, Illinois	Professor W. H. Marlow Logistics Research Project The Geo. Wash. University 707 - 22nd Street, N. W. Washington 7, D. C.
ASTIA Document Service Center Arlington Hall Station Arlington 12, Virginia	Professor L. W. Cohen Math. Dept., Univ. of Maryland College Park, Maryland	Professor Oskar Morgenstern Economics Research Project Princeton University 92 A Nassau Street Princeton, New Jersey
Office of Technical Services Department of Commerce Washington 25, D. C.	Professor Donald Eckman Director, Systems Research Center, Case Inst. of Tech. Cleveland, Ohio	Professor R. Radner Department of Economics University of California Berkeley, California
Technical Information Officer Naval Research Laboratory Washington 25, D. C.	Professor Lawrence E. Fouraker Dept. of Economics, The Pennsylvania State University State College, Pennsylvania	Professor Stanley Reiter Department of Economics Purdue University Lafayette, Indiana
C.O., ONR Branch Office 346 Broadway, New York 13, N. Y. Attn: J. Laderman	Professor David Gale Dept. of Math., Brown University Providence 12, Rhode Island	Mr. J. R. Simpson, Bureau of Supplies and Accounts (Code W31) Navy Department Washington 25, D. C.
C.O., ONR Branch Office 1030 East Green Street Pasadena 1, California Attn: Dr. A. R. Laufer	Professor L. Hurwicz School of Business Administration University of Minnesota Minneapolis 14, Minnesota	Professor A. W. Tucker Dept. of Mathematics Princeton University Princeton, New Jersey
Professor Russell Ackoff Operations Research Group Case Institute of Technology Cleveland 6, Ohio	Professor James R. Jackson Management Sciences Research Project, Univ. of California Los Angeles 24, California	Professor J. Wolfowitz Dept. of Mathematics Lincoln Hall, Cornell Univ. Ithaca 1, New York
Professor Kenneth J. Arrow Serra House Stanford University Stanford, California	Professor Samuel Karlin Dept. of Math., Stanford Univ. Stanford, California	